




Administering Glucagon

Call 911 immediately upon administration



1. Position the student on his or/her side.
2. Remove the cap from the glass vial.
3. Pull the needle cover off the syringe.
4. Insert the needle into vial and inject liquid.
5. Shake to dissolve.
6. Withdraw the glucagon solution back into the syringe and remove needle from vial.
7. Check for air bubbles in the syringe. Tap any visible air to the top of the syringe and gently push on the plunger until the air is removed.
8. Insert the needle at a 90-degree angle and inject the glucagon into a large muscle (upper arm, thigh, or upper outer area of buttock).
9. Withdraw the needle and apply slight pressure to the injection site.
10. Keep the student positioned on his other side.
11. Remain with the student until Emergency Medical Services (EMS) assumes control.

BAQSIMI GLUCAGON ADMINISTRATION



STEPS OF ADMINISTRATION

1. Verify that the student is unconscious or unable to swallow.
2. Take Baqsimi out of the container.
3. Hold device between fingers and thumb. **DO NOT** push plunger yet.




4. Insert tip gently into one nostril until fingers touch the outside of the nose.



5. Push plunger firmly all the way in. Dose is complete when the green line disappears.
6. Call 911 or have someone else call 911 immediately.
7. Stay with student until EMS arrives.
8. **Document Baqsimi glucagon administration using CCSD Medication Administration Log.**

GVOKE GLUCAGON ADMINISTRATION



1. Verify that the student is unconscious or unable to swallow.
2. Call 911 or have someone else call 911 immediately.
3. Tear open pouch at dotted line.
4. Remove syringe
5. Choose injection site and expose skin. Lower abdomen, outer thighs and outer upper arm are all appropriate sites.
6. Pull needle off cap. **DO NOT REMOVE AIR BUBBLES!**
7. **Pinch the skin and insert the needle at a 90° angle.**
8. **Push plunger down to inject glucagon.**
9. **Turn patient on side.**

Gvoke HypoPen (glucagon injection) **1 mg Auto-Inj**

1. Prepare → **2. Inject** → **3. Ass**

Tear Open Pouch at Dotted Line. Remove Auto-Injector.

Pull off Red Cap.

Choose Injection Site and Expose Skin.

Front View Back View

Lower Abdomen, Outer Thigh, or Outer Upper Arm

Push Down on Skin to Start. Hold Down for 5 Seconds. Wait for Window to Turn Red.

Hold Down for 5 Sec.

Turn Patient on Call Emergency

After the Injection, Put the Used Pen in a Safe Place Until It Can Be Disposed of Into a FDA Cleared Sharps Container.

Red Cap **Needle End**

10. Stay with student until EMS arrives.
11. Document Gvoke glucagon administration using CCSD Medication Administration Log.